

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 37/12	A1	(11) International Publication Number: WO 99/08521 (43) International Publication Date: 25 February 1999 (25.02.99)
(21) International Application Number: PCT/US98/16732 (22) International Filing Date: 13 August 1998 (13.08.98) (30) Priority Data: 60/055,614 14 August 1997 (14.08.97) US 60/055,613 14 August 1997 (14.08.97) US (71) Applicant: DEPARTMENT OF THE ARMY, U.S. GOVERNMENT (US/US); U.S. Army Medical Research & Material Command, 504 Scott Street, Fort Detrick, MD 21702-5012 (US). (72) Inventors: MEYERHOFF, James; 2936 Woodstock Avenue, Silver Spring, MD 20910 (US). HACKER, Henry; 9118 Jones Mill Road, Chevy Chase, MD 20815 (US). LONG, Joseph; 13833 Lakeside Drive, Clarksville, MD 21029 (US). (74) Agent: HENDRICKS, Glenna; P.O. Box 2509, Fairfax, VA 22031-2509 (US).		(81) Designated States: CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: TREATMENT OR PROPHYLAXIS OF RETINAL PATHOLOGY AND SPINAL CORD INJURY (57) Abstract Compositions containing 2-(phosphonomethyl)-pentanedioic acid (PMPA) and α and β N-acetyl-aspartyl-glutamate (NAAG) are appropriate means for treatment of neuronal tissue injury that may result from such factors as injury, toxicity, hypoxia or ischemia (a frequent cause of hypoxia). Target tissues include the brain, spinal cord and retina.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Title: Treatment or Prophylaxis of Retinal Pathology and Spinal Cord Injury

5 Field of the Invention:

This invention relates to the use of 2-(phosphonomethyl)-pentanedioic acid (PMPA) and α and β N-acetyl-aspartyl-glutamate (NAAG) for treatment of neuronal tissue injury that may result from such factors as injury, toxicity, hypoxia or ischemia (a frequent cause of hypoxia). Target tissues include the brain, spinal cord and retina.

10 Background of the Invention:

It is known that the neurotransmitter, N-Acetyl-aspartyl-glutamate (NAAG), which is found throughout the nervous system, is enzymatically converted to the potentially excitotoxic neurotransmitter, glutamate, by the enzyme α -linked acidic dipeptidase (NAALADase), also known as NAAG-hydrolyzing enzyme, in both normal and in pathophysiological conditions of the neurological system. (A new name was recently proposed for NAALADase: glutamate carboxypeptidase II.) Under current practice, drugs are used to block glutamate at its receptor "downstream". However, there has not previously been an adequate method for blocking the enzymatic formation of glutamate from an immediate precursor, NAAG.

25 It is believed that NAAG is released in response to neuronal trauma/ischemic injury, and is then hydrolyzed to form glutamic acid. Because NAAG has only a fraction of the excitatory potency of glutamic acid, the concentration of NAAG required to be neurotoxic might be 100-fold the toxic concentration of glutamic acid.

30 At present, spinal injury, whether arising from trauma or disease, disables many Americans of all ages. There has previously been no really effective therapeutic agent for use in treatment of such CNS injuries. Injuries such as those arising from penetrating injuries, exposure to blast, blunt trauma, falls and vehicular accidents as well as spinal cord and brain injury secondary to decompression sickness are

examples of instances when use of PMPA, α -NAAG or β -NAAG in accord with the teachings of this disclosure would be appropriate. Any damage which results in hypoxia, including ischemia and toxicity, can cause severe damage to the neuronal tissue.

5 A neuronal tissue that is particularly subject to damage arising from hypoxia, ischemia, exposure to toxins or any other causes of degeneration is the retina.

The leading causes of blindness in the United States include age-related macular degeneration (AMD), diabetic retinopathy, glaucoma and cataracts. The prevalence of
10 diabetes in North America is expected to reach almost 17 million by the year 2000. In cases of insulin-dependent diabetes mellitus (IDDM) with onset before age 30, the average prevalence of proliferative retinopathy is estimated at 23%.

15 In the case of IDDM of more than 30 years duration, however, the incidence of proliferative retinopathy rises to 70%. Proliferative retinopathy is the leading cause of new cases of blindness in the U.S., accounting for 12% of new cases annually.

20 The prevalence of AMD increases from over 2% in the age group 60-64 years to over 25% in the 75-80 year age group. It is estimated that the prevalence of glaucoma in the United States will be 2.9 million by the year 2000, and that over 130,000 will have been blinded by this disease.

25 At present, eye drops are available for treating vernal conjunctivitis containing both α - and β -NAAG. β -NAAG is a very weak NAALADase inhibitor that also has been found to act as an agonist. (α NAAG is also an agonist.) Thus, β -NAAG is not a specific NAALADase inhibitor. However, neither α -NAAG nor
30 β -NAAG were previously known to have use in treating or preventing damage to the retina or other neuronal tissue.

Summary of the Invention:

35 The instant invention is related to use of PMPA and α -NAAG and β -NAAG to prevent and treat conditions arising from exposure of neuronal tissue to toxins, injury, ischemia and hypoxia.

Detailed Description of the Invention:

It is now found that it is possible to inhibit the enzyme NAALADase, thereby blocking enzymatic conversion of NAAG to glutamate using PMPA. This approach is useful because blocking a metabolic source of glutamate may require less drug. Furthermore, previously known glutamate receptor blockers have unacceptable side effects.

The value of NAAG on the CNS was also studied. Hydrolysis of NAAG by N-acetylated α -linked acidic dipeptase (NAALADase) liberates the more potent excitatory amino acid neurotransmitter, glutamate (GLU). Consequently, NAAG has been regarded as a storage form of synaptic GLU and NAALADase has been proposed to modulate neuronal excitability by regulating the synaptic availability of GLU. NAAG has also been proposed to inhibit GLU release through activation of presynaptic mGluR3 receptors, and at low concentrations it has been shown to antagonize NMDA receptor-mediated responses.

It was found that it is possible to prevent toxicity or treat damage to neuronal tissue, including the spinal cord. Such damage may, for example arise from hypoxia in the CNS. Hence, traumatic injuries and diseases may be treated using PMPA and analogues which inhibit NAALADase may be used for treatment of such diseases. Injuries such as those arising from penetrating injuries, exposure to blast, blunt trauma, falls and vehicular accidents as well as spinal cord and brain injury secondary to decompression sickness are examples of instances when use of PMPA, α -NAAG and β -NAAG in accord with the teachings of this disclosure would be appropriate.

Studies in culture showed that PMPA protected spinal neurons from hypoxic injury in vitro. Furthermore, in studying the effect of PMPA, lumbar subarachnoid injections of dynorphin A (DYN) were administered to rats causing ischemic injury and paralysis that is mediated through excitatory amino acid receptors. The DYN was co-administered with PMPA at dosage of 0.4, 0.8 and 4 μ m. It was found that animals who received PMPA with the DYN showed evidence of significant protection from dynorphin-induced ischemia.

In order to study the effect of NAAG on the CNS, an accepted injury model was used to evaluate the pharmacological effects of NAAG (0.25-4 μ moles), specifically examining its effect on recovery of the persistent HL motor deficits elicited by L4-L5 subarachnoid injection of 10 η moles of dynorphine A (DYN). Spinal subarachnoid injections of NAAG alone did not alter HL motor function. When co-administered with DYN, NAAG caused significant dose-dependent improvements in motor scores by 24 hours post-injection. ($p < 0.05$, Kruskal-Wallis). Paralysis was still acutely evident after NAAG co-treatment. However, NAAG-induced improvements in recovery became increasingly apparent between 2 and 24 hours post-injection. The non-hydrolyzable analog β -NAAG also significantly improved recovery of HL motor function, indicating that this effect was not secondary to cleavage of NAAG into N-acetylaspartate and GLU.

The results indicate that NAAG has protective effects in instances of excitotoxic spinal cord injury that are likely to be mediated through NMDA receptors. Another form of injury for which prophylactic treatment is needed is occasioned by exposure of neuronal tissue to irradiation such as that which occurs in the spinal cord when irradiation is targeted for treatment of malignancies of the abdomen or in the retina when whole-head irradiation is administered for malignancies of the brain.

Dosage of NAAG and PMPA are essentially the same, and will, of course, depend on the age, size and condition of the patient as well as on the mode of administration.

For purposes of obtaining beneficial effects on the CNS, the agents of the invention can be delivered by any means which allows the active agent to contact the injured tissue. In the case of the spinal cord, the agents may be administered intrathecally or by caudal drip. Carriers which are appropriate are those which are basically non-irritating such as buffered saline which may contain glucose and other active agents such as antibiotics and anti-inflammatory agents. The concentration of PMPA is usually .05% to 6%. For example, a solution containing .5% PMPA in 5% glucose in phosphate

buffered saline may be administered intrathecally. Compositions containing PMPA may also be administered into the cisterna magna.

5 In addition to providing benefit for protection of the brain and spinal cord, PMPA, α -NAAG and β -NAAG may be used to prevent toxic damage to the retina, including damage arising from hypoxia in the retina. Hence, diseases which may be treated using PMPA and analogues which inhibit NAALADase may be used for treatment of such diseases as glaucoma, damage from
10 exposure to laser, blast over-pressure, blunt or penetrating injuries to the eye and photocoagulation such as that used in treatment of macular degeneration or diabetic retinopathy. Other diseases which affect the retinal vasculature such as lupus erythematosus and temporal arteritis are also responsible
15 for loss of sight. It appears a therapeutic window exists during the disease process when agents that reduce glutamate activity at its receptor sites can rescue neurons from injury.

Exposure of the retina to laser energy, whether therapeutic or accidental, results in formation of scotoma and visual
20 impairment. Even after therapeutic exposures, there may be immediate and progressive visual impairment due to destruction of normal retinal cellular elements with subsequent spread of injury to adjacent retinal tissue. Compositions containing PMPA may be administered prophylactically before expected
25 exposure to laser treatment or immediately after accidental exposure.

The agents of the invention can be delivered by any means which allows the active agent to contact the retina, including intraocular injection and administration of eye drops containing
30 carriers such as polyvinyl alcohol. Carriers which are appropriate are those which are basically non-irritating such as buffered saline containing 5% polyvinyl alcohol of such weight (number) as to be soluble in water. The PMPA may also be provided as a salve or ointment. (Higher concentrations of
35 polyvinyl alcohol will be useful in preparation of ointments.) The concentration of PMPA is usually .05% to 6%. For example, a solution containing 0.5% PMPA in phosphate buffered saline

containing 5% polyvinyl alcohol may be administered as drops at about 1-5 drops per dosing.

5 It has now been found that α -NAAG and β -NAAG also provide neuroprotective effects. Without relying on any particular theory for novelty, it is believed that these beneficial effects may result from competition for the glutamate receptor by α - and β -NAAG acting as a partial agonist at the N-methyl, D-aspartate (NMDA) receptor complex. It is also possible that the α -NAAG and β -Naag are acting as antagonists at the presynaptic mGluR3.

10 To model ischemias, sodium cyanide (NaCN) to inhibit oxidative metabolism and 2-deoxyglucose (2-DG) to inhibit glycolysis were used. Dissociated rabbit retina cells were studied to evaluate the potential neuroprotective effects of α - and β -NAAG. It was found that metabolic inhibition with NaCN/2-DG for 1 hour caused 50% toxicity when viability was assessed using the tetrazolium dye assay for succinate dehydrogenase activity. Co-treatment with α -NAAG resulted in dose-dependent protection of up to 55% ($p < 0.0005$, T-test). 15 When the unhydrolysable β -NAAG was employed, a dose-dependent protection of up to 37% was observed ($p < 0.001$, t-test).

20 The active agents of the invention are readily available commercially. Compositions containing PMPA and α - and β -NAAG may be administered for treatment of conditions of the retina by such means as eye drops or by injection. Appropriate carriers include saline, glucose (5% or 10% being more usual) in half normal saline, and buffered saline, etc. Compositions may also contain other agents used as carriers such as dimethylsulfoxide (DMSO).

25 For intraocular administration, as an example, a 0.05% solution of PMPA in saline may be injected into the eye.

30 Compositions for use in the method of this invention may contain, in addition to PMPA, other active agents such as anti-inflammatory agents, antibiotics, alkaloids, anesthetics and analgesics.

35 The dosage administered will depend on the condition, size and age of the patient. In general, dosage of 0.0001/Kg to 1

mg/Kg for PMPA, α -NAAG or β -NAAG will be appropriate, with larger animals such as man receiving about 0.0001 to 0.05 mg/Kg and smaller animals receiving larger dosage relative to weight.

5 Compositions for use in the method of this invention may contain, in addition to PMPA, other active agents such as anti-inflammatory agents, antibiotics, anesthetics and analgesics.

Chronic depot delivery systems known in the art may be used to deliver the active agents. For purposes of treating retinal diseases, the active agents may also be administered
10 intraocularly such as, for example, by intravitreal means.

Example 1:

A composition for administration by injection (including intrathecal injection) is prepared by adding 3 mg PMPA to 3 ml DMSO. To this is added sufficient 5% glucose in half-normal
15 saline to make 100 ml solution.

Example 2:

A composition is prepared for use as eye drops. Six mg of β -NAAG is added to 3 ml DMSO. To this is then added sufficient saline to make 100 ml of solution.

20 **Example 3:**

A composition is prepared for intrathecal injection by adding sufficient 10% glucose in half-normal saline to 3 mg of α -NAAG to provide 100 ml of solution.

It is urged that the above examples are provided for
25 purposes of general instruction and are not to be construed as in any way limiting the scope of the invention.

What we claim is:

1. A composition of matter comprising 2-(phosphonomethyl) pentanedioic acid or one of α and β N-acetyl-aspartyl-glutamate (NAAG) in a pharmaceutically acceptable carrier in a neuronal protective effective amount.
5
2. A composition of claim 1 wherein the 2-(phosphonomethyl) pentanedioic is present in a concentration of .05% to 6%.
10
3. A method of treating or protecting against neurological damage comprising administration of a composition of matter comprising a glutamate formation inhibiting effective amount of 2-(phosphonomethyl) pentanedioic acid, α -NAAG or β -NAAG in a pharmaceutically acceptable carrier.
15
4. A method of claim 3 wherein the composition contains 2-(phosphonomethyl) pentanedioic and is administered intrathecally.
20
5. A method of claim 3 wherein the neurological damage arises from trauma, vertebral disease or decompression sickness.
25
6. A method of claim 3 wherein the neuronal tissue requiring treatment or protection is retinal tissue.
7. A method of claim 6 wherein the composition is administered in the form of eye drops.
30
8. A method of claim 3 wherein the composition contains 0.5% to 6% PMPA, α -NAAG or β -NAAG.
- 35 9. A method of claim 8 wherein the active agent is PMPA.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/16732

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A01N 37/12

US CL : 514/121, 563

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/121, 563

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONEElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,000,936 A (CHIBRET) 19 March 1991, see abstract.	1 and 2
Y	Database HCAPLUS on STN, No. 1996:6894, JACKSON et al., "Design, Synthesis, and Biological Activity of a Potent Inhibitor of the Neuropeptidase An-Acetylated .alpha.-linked Acidic Dipeptidase", abstract to J. Med. Chem. 39(2), 619-622, 1996.	1-9



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

B earlier document published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

A

document member of the same patent family

Date of the actual completion of the international search

15 NOVEMBER 1998

Date of mailing of the international search report

23 DEC 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DWAYNE C. JONES

Telephone No. (703) 308-1235

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/16732

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

REGISTRY, HCAOLD, HCAPLUS, USPATFULL, EMBASE, BIOSIS, WPIDS structure search with the following terms: phosphonomehtylpentanedioic or phosphonomethyl, L-glutamic acid, n-acetyl-d-beta-aspartyl, NAAAG, acetylaspartylglutamate or acetylaspartylglutamic